

Stewardship applied to antipsychotics: Development of an antipsychotic stewardship program in inpatient settings for monitoring and optimizing outcomes

Gillian Gonzales, PharmD¹

Kari Tornes, PharmD²

Stephen R. Saklad, PharmD, BCPP³

How to cite: Gonzales G, Tornes K, Saklad SR. Stewardship applied to antipsychotics: development of an antipsychotic stewardship program in inpatient settings for monitoring and optimizing outcomes. *Ment Health Clin* [Internet]. 2022;12(5):320-6. DOI: 10.9740/mhc.2022.10.320.

Submitted for Publication: September 28, 2021; **Accepted for Publication:** August 2, 2022

Abstract

Antipsychotic (AP) medications are prescribed for various psychiatric diagnoses that require routine monitoring to ensure optimal use, effectiveness, adherence, and for potentially severe adverse effects. There is currently no comprehensive protocol for institutional supervision of prescribing and monitoring AP. Antibiotics (ABX) are commonly associated with stewardship programs aimed at optimizing use and mitigating harm. These programs have proven to result in positive outcomes in both safety and efficacy parameters for numerous institutions. Given that AP are also associated with significant adverse effects and often misused, the concept of stewardship can be applied to this class of agents to optimize their use and improve overall patient outcomes. The objective of this paper is to provide guidance for the implementation of antipsychotic stewardship programs (APSP) in the inpatient setting. The development of this APSP was designed based on ABX stewardship programs and the Centers for Disease Control and Prevention, Agency for Healthcare Research and Quality, and the American Psychiatric Association practice guidelines on the treatment of patients with schizophrenia. In conclusion, APSPs have the potential to enhance and standardize institutional supervision of prescribing and monitoring practices of AP, leading to improved clinical outcomes and the reduction of adverse effects. APSP teams should be multidisciplinary, consisting of clinicians and administrators, working in conjunction with patients and patient advocates to design individualized recovery plans that consider the individual patient's history and desired outcomes. Monitoring, stewardship interventions, and outcomes should be documented on both an individual and deidentified institutional basis, analyzed, and summarized periodically as measures for quality improvement.

Keywords: antipsychotics, stewardship, monitoring, quality improvement, schizophrenia, bipolar disorder, major depressive disorder, agitation, behavioral disturbance, response, adverse effects

¹ PGY1 Resident, VA Eastern Colorado Health Care System, Aurora, Colorado; previously: The University of Texas at Austin College of Pharmacy, San Antonio, Texas, ORCID: <https://orcid.org/0000-0002-8072-9744>; ² PGY1 Resident, University Health System, San Antonio, Texas; previously: The University of Texas at Austin College of Pharmacy, San Antonio, Texas, ORCID: <https://orcid.org/0000-0002-8605-4688>; ³ (Corresponding author) Clinical Professor and Director of Psychiatric Pharmacy, Pharmacotherapy Division, College of Pharmacy, University of Texas at Austin, San Antonio, Texas, Saklad@uthscsa.edu, ORCID: <https://orcid.org/0000-0002-6181-138X>

Disclosures: GG and KT have nothing to disclose. SRS is an employee of The University of Texas at Austin College of Pharmacy; appointed to the

Texas Health and Human Services Commission, San Antonio State Hospital, and the UT Health San Antonio Long School of Medicine; is a consultant for Alkermes, BioXcel, Intra-Cellular Therapies, Genomind, Janssen, Karuna, and Lyndra; is a part of the speakers bureau for Intra-Cellular Therapies, Otsuka PsychU, Neurocrine, Teva, Texas Society of Health-System Pharmacists; and is an occasional speaker for several professional organizations; additionally, he is a part of the Business Development Council for the College of Psychiatric and Neurologic Pharmacists and, last, serves as an expert witness on both defendant and plaintiff sides.



© 2022 AAPP. The *Mental Health Clinician* is a publication of the *American Association of Psychiatric Pharmacists*. This is an open access article distributed under the terms of the *Creative Commons Attribution-NonCommercial 3.0 License*, which permits non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

TABLE 1: Warnings and precautions for antipsychotics use as outlined by FDA-approved product labels

Stroke (cerebrovascular problems) in elderly people that can lead to death.

Neuroleptic malignant syndrome: high fever; confusion; changes in breathing, heart rate, and blood pressure; stiff muscles and increased sweating; these may be symptoms of a rare but potentially fatal condition.

Uncontrolled body movements (tardive dyskinesia) of the face, tongue, or other body parts.

Problems with metabolism, including high blood sugar, diabetes, increased fat (cholesterol and triglyceride) levels in the blood, and weight gain.

Low white blood cell count

Decreased blood pressure (orthostatic hypotension). Feeling lightheaded, dizzy, or faint when quickly arising from a sitting or lying position.

Falls. Feeling sleepy or dizzy due to a decrease in blood pressure when changing position (orthostatic hypotension). Additionally, can slow thinking and motor skills, which may lead to falls that can result in broken bones or other injuries.

Seizures (convulsions).

Sleepiness and trouble concentrating: Do not drive, operate heavy machinery, or do other dangerous activities until the effect of the antipsychotic is known.

Problems regulating increased body temperature. Avoid getting overheated or dehydrated while taking the antipsychotic.

Difficulty swallowing.

Introduction

Psychotropic agents have transformed the care of patients with mental disorders over the past 70 years.¹ Antipsychotic (AP) agents are now indicated and frequently prescribed to patients with various psychiatric diagnoses, including schizophrenia and bipolar disorder and as an adjunct in major depressive disorder. Despite this success, AP are found to be associated with problematic and potentially severe adverse effects. There are no trivial indications for AP, and all use requires careful patient selection following comprehensive baseline assessment, medication choice, medication starting dose and titration, and ongoing monitoring. There currently is no comprehensive framework for systematic supervision of prescribing and monitoring practices of AP to enhance patient safety and efficacy. The current system of AP monitoring is primarily based on each individual clinician's interpretation and compliance with published guidelines and facility policies. As shown in Table 1, the FDA-approved product labels for AP contain warnings and precautions of which patients should be aware and discuss with their health care providers. Adverse effects of AP use include weight gain, sedation, metabolic syndrome, glucose-insulin homeostasis disruption, prolactin elevation, drug-induced parkinsonism, and tardive dyskinesia.^{2,3} Dosage adjustments and monitoring for the early appearance of adverse effects can reduce discomfort and the risk for more severe adverse effects to become apparent.³

A comprehensive monitoring program, such as an antipsychotic stewardship program (APSP), formalizes the patient-monitoring processes that already exist in various forms and creates a framework that can be successfully replicated in many clinical care settings. Suboptimal AP use contributes to the risk of adverse effects and harmful long-term health consequences, potentially without adequate control of

psychotic symptoms.⁴ Cardiovascular conditions remain a key cause of death among people treated with AP, suggesting suboptimal monitoring practices delay the identification of high-risk patients and prevent patients from receiving proper management.^{5,6} Studies⁶⁻⁹ show that rates of metabolic monitoring are inadequate and there are significant variations in the monitoring practices of cardiometabolic outcomes in people who use AP, further demonstrating the need for standardized monitoring practices.

To optimize efficacy and reduce harms associated with AP use, AP should be routinely prescribed and monitored following best practices. This situation parallels the use of antibiotics (ABX) with the potential for profound benefits and avoiding serious adverse effects when used appropriately. The Centers for Disease Control and Prevention (CDC)¹⁰ estimates that about 30% of all ABX prescribed in acute settings in the United States are suboptimal or unnecessary. ABX stewardship programs were developed to improve ABX prescribing and monitoring, leading to improved outcomes with minimal harms to patients.¹¹

ABX stewardship is a systematic and multidisciplinary approach aimed at optimization and appropriate use of ABX to promote the best clinical outcomes for patients, avoid adverse consequences, and limit the development of ABX resistance.^{12,13} Studies¹²⁻¹⁴ show that implementation of ABX stewardship initiatives results in positive outcomes, including a relative risk reduction in mortality associated with guideline-adherent empirical therapy and de-escalation to more targeted therapy after cultures are identified. The concept of stewardship, developed initially to optimize the use of ABX, can be successfully applied to AP use with the goal of encouraging optimal use and monitoring to improve efficacy, safety, and tolerability. A clinician-driven inter-

vention that evaluates the patient's comprehensive medical history, previous medication use, indication for an AP, adverse effect monitoring, and rationale for continuing medications long term can improve outcomes and reduce the adverse effects in patients.² We provide guidance to develop an APSP, including essential team members and suggestions for implementation.

Methods

A search for meta-analyses and systematic reviews of ABX stewardship programs was conducted on PubMed without restrictions and used to identify the impact of ABX stewardship program interventions.¹⁴⁻¹⁸ The CDC Core Elements of Hospital Antibiotic Stewardship Programs¹⁰ and the Agency for Healthcare Research and Quality¹³ Acute Care Hospital Toolkit to Improve Antibiotic Use were employed as prototypes to determine key components that were found to be necessary for the development of a successful ABX stewardship program.

This guidance for APSP implementation in the inpatient setting was based on the experience and recommendations implementing a successful ABX stewardship program in a similar setting. Among all types of psychotropics, it was decided that antipsychotics would be the first agents around which to develop a stewardship program as they have the most potential harms if not prescribed and monitored following best practices. There are some unique differences that require adaptation. For example, mental health clinics have most of their patients seen over an extended period with many interruptions in continuity. The American Psychiatric Association Practice Guidelines on the Treatment of Patients with Schizophrenia have recently been revised and can currently be used to inform individual APSPs algorithms for patient selection, assessment, and monitoring during treatment with AP.¹⁹

APSP Core Elements

APSP can incorporate the core elements of ABX stewardship as described by the CDC, which may facilitate ease of implementation in health care facilities that have already established an ABX stewardship program due to familiarity of concepts.¹⁰ Those health care facilities without a preexisting ABX stewardship program require additional educational components to introduce the seven core elements of the proposed APSP. These core elements include the following:

- Institutional leadership commitment
- Accountability
- Pharmacist expertise
- Action
- Tracking

- Reporting
- Education¹⁰

Institutional Leadership Commitment

The buy-in and involvement from senior mental health facility executives is essential to further the goals of the program. The facility administrators need to identify and dedicate the necessary human, financial, and information technology resources that are needed to implement and maintain the APSP. Institutional leadership is responsible for connections with stakeholders and demonstrating that the APSP is supported by the healthcare facility.

Accountability

APSP should have explicitly identified APSP coleaders (we recommend a psychiatrist and pharmacist) that have the authority and resources and are responsible for its outcomes. APSP leaders must have the necessary skills, including leadership, project management, communication, and training in psychiatry and psychopharmacology. Responsibilities and expectations should be clear and explicit to provide accountability for the program.

Pharmacist Expertise

Engagement from pharmacy is essential for ensuring the effectiveness of the APSP. The pharmacy department should identify an experienced pharmacist as coleader of the facility APSP. Pharmacists trained and board certified in the treatment of psychiatric conditions are effective in ensuring the best practices in the use of APs.²⁰ A board-certified psychiatric pharmacist can serve the organization as a specialist through clinical decision making; implementing best practices; and educating health care personnel, patients, and the public.

Action

The primary APSP intervention is prospective review of AP orders and providing the prescriber feedback to follow best practices as implemented by the facility. AP use for the most common indications and first-line drug choice are routine and should require few prospective interventions. AP monitoring for treatment of emergent problems with adverse effects, efficacy, and adherence are opportunities for increased APSP interventions. Confirming baseline and ongoing documentation should be automated to the greatest extent possible and provide prospective alerts for required time- and event-based assessments.

Tracking

Quarterly reviews of the stewardship program should be performed as a quality assurance measure. See Figure 1. These reviews should include the confirmation of baseline documentation of the rationale to use a specific AP, informed consent including disclosure of potential risks and possible benefits, consideration of the patient's health care beliefs in the recovery plan, and the use of shared decision making to support the appropriateness of the prescribed medication.²¹ Ongoing monitoring of APs after initiation should include starting dose and titration as well as the reason for subsequent AP adjustments and their




FIGURE 1: Antipsychotic stewardship patient care process

outcomes.^{19,22} The rationale for the AP changes may include patient response, adverse effects, dose changes that were planned as part of the titration, added medications, discontinued medications, and changes in route or timing of medications.

Reporting

These reviews should be summarized, analyzed, and reported quarterly at the facility level. More frequent reports would be made as appropriate on an individual patient basis. The intent of the tracking and reports on a facility basis is to provide insight on what is working well and identify potential areas for improvement. The individual patient reporting is intended to ensure best practices are being followed. Criteria for medication use should be monitored in the literature and adjusted periodically to ensure that best practices are being followed. Revisions based on reported outcomes should be implemented following the plan-do-study-act or equivalent problem-

solving model.^{23,24} Figure 2 depicts the ongoing quality assurance process necessary for a successful program.

Education

APSP should incorporate a routine education component to review best practices incorporating new evidence on existing drugs, new drugs and their place in treatment, and summarized findings from the quarterly reports. Education programs must include customized components tailored for the unique needs of clinicians, administrators, and stakeholders to encourage continuity of the stewardship initiative.

APSP Team Composition

Overall, clinicians are vital collaborators in stewardship development. Clinician buy-in and support is essential to ensure that best practice guidelines are adopted and adhered to so that the program is successful. See Table 2.

Pharmacist

Pharmacists can support the APSP team by conducting interventions, coordinating data needs, ensuring appropriate use of medications, and serving as a bridge to the pharmacy department. The pharmacist should have board certification in psychiatric pharmacy and the ability to communicate and advise physicians, other clinicians, administrators, and patients in AP use best practices.

Psychiatrist

The psychiatrist should collaborate with the pharmacist to decide the goals of the program. The psychiatrist should see and evaluate patients for clarification or confirmation of diagnoses. Additionally, the psychiatrist should have oversight of prescribing recommendations and settle differences of opinions between the APSP and other prescribers. The psychiatrist should have board certification




FIGURE 2: Antipsychotic stewardship system quality assurance process

TABLE 2: Clinicians and administrators necessary for successful implementation of an APSP

Pharmacists
Physicians
Social workers
Nurses
Patients
Pharmacy and therapeutics (liaison)
Information technology (liaison)
Quality improvement
Other supervisory or regulatory bodies (liaison)

in psychiatry and the ability to communicate and advise physicians, other clinicians, administrators, and patients in AP use best practices.

Clinical Social Worker

The clinical social worker provides case management services, individual and group therapy, conducts biopsychosocial assessments and social histories, determines accessibility and patient eligibility for services (including collaboration on access to specialized medication options, such as long-acting injectable AP), assists in coordination and transitions of care, and develops recovery and discharge plans. The clinical social worker should have training in psychiatric conditions and ability to communicate plans with other team members.

Pharmacy Department

The pharmacy department is responsible for processing medication orders and ensuring medications are delivered promptly. Pharmacy can help ensure that prescribing protocols are followed, assist with drug-monitoring protocols, and field drug-related questions. Additionally, pharmacy can collect data on AP use, observe trends in prescribing and dispensing of AP medications, and disseminate information to other clinicians.

Nursing

Nursing is responsible for helping assess patient status, response to interventions, and monitoring for adverse effects. Due to increased direct interaction with patients, nursing is a vital component of the APSP. Nursing preferably will have specialized training in psychiatry and the ability to communicate and advocate for patient needs to both treatment team members and the APSP.

Administrative Support

Administrative support for an APSP is required for the successful implementation and longevity of the program. Suggestions for roles and contributions of various administrators are outlined below.

Executive Leadership

Involvement of executive leadership promotes and supports the goals of the stewardship program. Leadership may also

assist in identifying and allocating necessary resources for the APSP.

Pharmacy and Therapeutics (P&T)

The P&T committee oversees the formulary and may endorse prescribing guidelines. The committee may also coordinate incorporation of APSP goals and guidelines with other committees.

Quality Improvement

Quality improvement can ensure compliance with standards and other regulations of regulatory bodies, ensure compliance with quality metrics while promoting rational AP use, and provide resources for interventions to improve AP use.

Information Technology (IT)

IT may aid with data collection and integration of stewardship protocols into operational systems.

Patient-Centered Care

Patient input and involvement is essential in developing a successful, goal-concordant recovery plan. Without patient engagement and buy-in, it is unlikely that an intervention will be successful. Multiple factors should be considered, including the patient's past medical history; history of present illness; social history, including alcohol, tobacco, and substance use; diet; and current and past medications, including all AP, other psychotropics, and all medications for nonpsychiatric disorders. Patient goals, values, and the complexity of the regimen should be considered as well. The health care belief model can be used to understand the patient's perceptions and motivations toward behavior and actions affecting diagnoses.²⁵ Effective communication is essential in ensuring that patients and providers can engage in shared decision making.²⁶ Providers should be nonjudgmental, using open-ended questions to elicit patient information, views, and goals and encourage involvement. It is important for patients to believe that they have a condition, that their condition can be treated, and that the benefits of the treatment outweigh any risks. Patients must understand their individual recovery plan, including medication and nonmedication treatment regimens, alternatives to the recommendations, and realistic expectations for the outcome if they participate and follow their recovery plan.

Discussion

AP medications are routinely used to treat psychiatric conditions but have significant adverse effects that require monitoring. The concept of stewardship can be applied to AP use in health care settings to promote safe use and optimize therapy. APSP teams should be multidisciplinary, consisting of clinicians and administrators working in conjunction with patients and patient advocates to design individualized recovery plans that consider the individual patient's history and desired outcomes. The risks and benefits and appropriateness of AP use should be assessed

and monitored routinely for each patient based on the patient's medical and social history, current status, and goals of therapy. Monitoring, stewardship interventions, and outcomes should be documented on both an individual and deidentified institutional basis and analyzed and summarized periodically as measures for quality improvement. APSPs have the potential to improve and standardize prescribing supervision, monitoring, and documentation, leading to improved outcomes of therapy. This document provides guidance for the development and implementation of APSPs in inpatient settings to monitor and optimize patient outcomes while receiving AP therapy.

References

- Lehmann HE, Ban TA. The history of the psychopharmacology of schizophrenia. *Can J Psychiatry*. 1997;42(2):152-62. DOI: [10.1177/070674379704200205](https://doi.org/10.1177/070674379704200205). PubMed PMID: [9067064](https://pubmed.ncbi.nlm.nih.gov/9067064/).
- Kovacic NL, Gagnon DJ, Riker RR, Wen S, Fraser GL. An analysis of psychoactive medications initiated in the ICU but continued beyond discharge: a pilot study of stewardship. *J Pharm Pract*. 2020;33(6):760-7. DOI: [10.1177/0897190019830518](https://doi.org/10.1177/0897190019830518). PubMed PMID: [30813837](https://pubmed.ncbi.nlm.nih.gov/30813837/).
- Meyer JM. Pharmacotherapy of Psychosis and Mania. In: Brunton LL, Hilal-Dandan R, Knollmann BC editors. *Goodman & Gilman's: the pharmacological basis of therapeutics*. 13th ed. New York: McGraw-Hill; 2017.
- Solmi M, Murru A, Pacchiarotti I, Undurraga J, Veronese N, Fornaro M, et al. Safety, tolerability, and risks associated with first- and second-generation antipsychotics: a state-of-the-art clinical review. *Ther Clin Risk Manag*. 2017;13:757-77. DOI: [10.2147/TCRM.S117321](https://doi.org/10.2147/TCRM.S117321). PubMed PMID: [28721057](https://pubmed.ncbi.nlm.nih.gov/28721057/); PubMed Central PMCID: [PMC5499790](https://pubmed.ncbi.nlm.nih.gov/PMC5499790/).
- Jakobs KM, Posthuma A, de Grauw WJC, Schalk BWM, Akkermans RP, Lucassen P, et al. Cardiovascular risk screening of patients with serious mental illness or use of antipsychotics in family practice. *BMC Fam Pract*. 2020;21(1):153. DOI: [10.1186/s12875-020-01225-7](https://doi.org/10.1186/s12875-020-01225-7). PubMed PMID: [32727372](https://pubmed.ncbi.nlm.nih.gov/32727372/); PubMed Central PMCID: [PMC7391510](https://pubmed.ncbi.nlm.nih.gov/PMC7391510/).
- Azfr Ali RS, Jalal Z, Paudyal V. Guidelines versus practice in screening and monitoring of cardiometabolic risks in patients taking antipsychotic medications: where do we stand? *Gen Psychiatr*. 2021;34(4):e100561. DOI: [10.1136/gpsych-2021-100561](https://doi.org/10.1136/gpsych-2021-100561). PubMed PMID: [34396043](https://pubmed.ncbi.nlm.nih.gov/34396043/); PubMed Central PMCID: [PMC8311327](https://pubmed.ncbi.nlm.nih.gov/PMC8311327/).
- Keenan R, Chepulis L, Ly J, Carter S, Lao C, Asim M, et al. Metabolic screening in primary care for patients with schizophrenia or schizoaffective disorder and taking antipsychotic medication. *J Prim Health Care*. 2020;12(1):29. DOI: [10.1071/HC19023](https://doi.org/10.1071/HC19023). PubMed PMID: [32223847](https://pubmed.ncbi.nlm.nih.gov/32223847/).
- Mitchell AJ, Delaffon V, Vancampfort D, Correll CU, De Hert M. Guideline concordant monitoring of metabolic risk in people treated with antipsychotic medication: systematic review and meta-analysis of screening practices. *Psychol Med*. 2012;42(1):125-47. DOI: [10.1017/S003329171100105X](https://doi.org/10.1017/S003329171100105X). PubMed PMID: [21846426](https://pubmed.ncbi.nlm.nih.gov/21846426/).
- Hibner TA, Wakefield AN, Eaves SM, Gonzalvo JD, Macik MR, Williams GD. Metabolic monitoring of second-generation antipsychotics: Evaluation of a pharmacist- and nurse-driven protocol. *J Am Pharm Assoc* (2003). 2020;60(5):S88-92. DOI: [10.1016/j.japh.2020.04.016](https://doi.org/10.1016/j.japh.2020.04.016). PubMed PMID: [32513507](https://pubmed.ncbi.nlm.nih.gov/32513507/).
- CDC. Core Elements of Hospital Antibiotic Stewardship Programs [cited 2020 Sep 14]. Atlanta: US Department of Health and Human Services, CDC; 2019. Available from <https://www.cdc.gov/antibiotic-use/core-elements/hospital.html>
- Dellit TH, Owens RC, McGowan Jr JE, Gerding DN, Weinstein RA, Burke JP, et al. Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America guidelines for developing an institutional program to enhance antimicrobial stewardship. *Clin Infect Dis*. 2007;44(2):159-77. DOI: [10.1086/510393](https://doi.org/10.1086/510393). PubMed PMID: [17173212](https://pubmed.ncbi.nlm.nih.gov/17173212/).
- Barlam TF, Cosgrove SE, Abbo LM, MacDougall C, Schuetz AN, Septimus EJ, et al. Implementing an antibiotic stewardship program: guidelines by the Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America. *Clin Infect Dis*. 2016;62(10):e51-77. DOI: [10.1093/cid/ciw118](https://doi.org/10.1093/cid/ciw118). PubMed PMID: [27080992](https://pubmed.ncbi.nlm.nih.gov/27080992/); PubMed Central PMCID: [PMC5006285](https://pubmed.ncbi.nlm.nih.gov/PMC5006285/).
- Agency for Healthcare Research and Quality [Internet]. Toolkit to improve antibiotic use in acute care hospitals [cited 2019 Nov]. Rockville (MD): Agency for Healthcare Research and Quality. Available from: <https://www.ahrq.gov/antibiotic-use/acute-care/index.html>
- Monnier AA, Schouten J, Le Maréchal M, Tebano G, Pulcini C, Stanić Benić M, et al. Quality indicators for responsible antibiotic use in the inpatient setting: a systematic review followed by an international multidisciplinary consensus procedure. *J Antimicrob Chemother*. 2018;73(suppl_6):vi30-9. DOI: [10.1093/jac/dky116](https://doi.org/10.1093/jac/dky116). PubMed PMID: [29878221](https://pubmed.ncbi.nlm.nih.gov/29878221/); PubMed Central PMCID: [PMC5989598](https://pubmed.ncbi.nlm.nih.gov/PMC5989598/).
- Schweitzer VA, van Werkhoven CH, Rodríguez Baño J, Bielicki J, Harbarth S, Hulscher M, et al. Optimizing design of research to evaluate antibiotic stewardship interventions: consensus recommendations of a multinational working group. *Clin Microbiol Infect*. 2020;26(1):41-50. DOI: [10.1016/j.cmi.2019.08.017](https://doi.org/10.1016/j.cmi.2019.08.017). PubMed PMID: [31493472](https://pubmed.ncbi.nlm.nih.gov/31493472/).
- Schuts EC, Hulscher MEJL, Mouton JW, Verduin CM, Stuart JWTC, Overdiek HWPM, et al. Current evidence on hospital antimicrobial stewardship objectives: a systematic review and meta-analysis. *Lancet Infect Dis*. 2016;16(7):847-56. DOI: [10.1016/S1473-3099\(16\)00065-7](https://doi.org/10.1016/S1473-3099(16)00065-7). PubMed PMID: [26947617](https://pubmed.ncbi.nlm.nih.gov/26947617/).
- Falcone M, Paul M, Yahav D, Orlando G, Tiseo G, Prendki V, et al. Antimicrobial consumption and impact of antimicrobial stewardship programmes in long-term care facilities. *Clin Microbiol Infect*. 2019;25(5):562-9. DOI: [10.1016/j.cmi.2018.07.028](https://doi.org/10.1016/j.cmi.2018.07.028). PubMed PMID: [30076978](https://pubmed.ncbi.nlm.nih.gov/30076978/).
- Le Maréchal M, Tebano G, Monnier AA, Adriaenssens N, Gysens IC, Huttner B, et al. Quality indicators assessing antibiotic use in the outpatient setting: a systematic review followed by an international multidisciplinary consensus procedure. *J Antimicrob Chemother*. 2018;73(suppl_6):vi40-9. DOI: [10.1093/jac/dky117](https://doi.org/10.1093/jac/dky117). PubMed PMID: [29878218](https://pubmed.ncbi.nlm.nih.gov/29878218/).
- Keepers GA, Fochtman LJ, Anzia JM, Benjamin S, Lyness JM, Mojtabei R, et al. The American Psychiatric Association Practice Guideline for the treatment of patients with schizophrenia. *Am J Psychiatry*. 2020;177(9):868-72. DOI: [10.1176/appi.ajp.2020.177901](https://doi.org/10.1176/appi.ajp.2020.177901). PubMed PMID: [32867516](https://pubmed.ncbi.nlm.nih.gov/32867516/).
- Goldstone LW, DiPaula BA, Caballero J, Park SH, Price C, Zasadzki Slater M. Improving medication-related outcomes for patients with psychiatric and neurologic disorders: value of psychiatric pharmacists as part of the health care team. *Mental Health Clin* [Internet]. 2015;5(1):1-28. DOI: [10.9740/mhc.2015.01.00](https://doi.org/10.9740/mhc.2015.01.00)
- ABIM Foundation. American Psychiatric Association: choosing wisely. Five things physicians and patients should question [cited 2019 Sep 20]. Available from: <https://www.choosingwisely.org/societies/american-psychiatric-association/>.
- American Diabetes Association, American Psychiatric Association, American Association of Clinical Endocrinologists, North American Association for the Study of Obesity. Consensus development conference on antipsychotic drugs and obesity and diabetes.

- Diabetes Care. 2004;27(2):596-601. DOI: [10.2337/diacare.27.2.596](https://doi.org/10.2337/diacare.27.2.596). PubMed PMID: [14747245](https://pubmed.ncbi.nlm.nih.gov/14747245/).
23. Agency for Healthcare Research and Quality [Internet]. Plan-do-study-act (PDSA) directions and examples [cited 2020 Sep]. Rockville (MD): Agency for Healthcare Research and Quality. Available from: <https://www.ahrq.gov/health-literacy/improve/precautions/tool2b.html>
 24. Institute for Healthcare Improvement [Internet]. How to improve [cited 2020 Sep 14]. Available from: <http://www.ihl.org/resources/Pages/HowtoImprove/default.aspx>
 25. Jones CL, Jensen JD, Scherr CL, Brown NR, Christy K, Weaver J. The Health Belief Model as an explanatory framework in communication research: exploring parallel, serial, and moderated mediation. *Health Commun.* 2015;30(6):566-76. DOI: [10.1080/10410236.2013.873363](https://doi.org/10.1080/10410236.2013.873363). PubMed PMID: [25010519](https://pubmed.ncbi.nlm.nih.gov/25010519/).
 26. Sanders JJ, Curtis JR, Tulsy JA. Achieving goal-concordant care: a conceptual model and approach to measuring serious illness communication and its impact. *J Palliat Med.* 2018;21(S2):S17-27. DOI: [10.1089/jpm.2017.0459](https://doi.org/10.1089/jpm.2017.0459). PubMed PMID: [29091522](https://pubmed.ncbi.nlm.nih.gov/29091522/).